



Phytochemical characterization and antioxidant profile of *Sechium edule* (Jacq) Swartz (*Cucurbitaceae*) varieties grown in Costa Rica

[Caracterización fitoquímica y perfil antioxidante de las variedades de *Sechium edule* (Jacq) Swartz (*Cucurbitaceae*) cultivadas en Costa Rica]


Jonathan Parra^{1*}, Priscilla Hernández¹, Fernando Ocampo-Maroto², Víctor Álvarez-Valverde², Yendry Carvajal-Miranda², Gerardo Rodríguez-Rodríguez², Cristina Herrera¹

¹Instituto de Investigaciones Farmacéuticas (INIFAR), Facultad de Farmacia, Universidad de Costa Rica, Sede Rodrigo Facio, San Pedro, San José, 11501-2060. Costa Rica.

²Laboratorio de Fitoquímica, Escuela de Química, Universidad Nacional de Costa Rica, Campus Omar Dengo, Heredia, 86-3000. Costa Rica.

*E-mail: jonathan.parra@ucr.ac.cr


SUPPLEMENTARY DATA




S1. Chromatogram for the HPLC separation of **1**. Retention time 11.38 min.




S2. Chromatogram for the HPLC separation of **2**. Retention time 9.79 min.




S3. ¹H-NMR spectra of 1.




S4. ^{13}C -NMR spectra of 1.




S5. HSQC spectra of 1.




S6. H-H COSY spectra of **1**.




S7. HMBC spectra of 1.




S8. ¹H-NMR spectra of 2.




S9. ¹³C-NMR spectra of 2.



S10. HSQC spectra of 2.



S11. H-H COSY spectra of 2.



S12. HMBC spectra of 2.