



Gallic acid protects against cadmium chloride-induced alterations in Wistar rats via the antioxidant defense mechanism

[El ácido gálico protege contra las alteraciones inducidas por el cloruro de cadmio en ratas Wistar a través del mecanismo de defensa antioxidante]

Oluwafemi Adeleke Ojo^{1,2*}, Damilare Rotimi², Ayomide Emmanuel Bright², Omowumi Titilola Kayode³, Adebola Busola Ojo⁴, Omokolade Oluwaseyi Alejlowo^{1,2}, Basiru Olaitan Ajiboye^{5,6}, Olarewaju M Oluba^{1,2}

¹Landmark University SDG 03 (Good Health and Well Being Research Group), Nigeria.

²Department of Biochemistry, Landmark University, Omu-Aran, Nigeria.

³Biochemistry Unit, Department of Biological Sciences, Mountain Top University, Prayercity, Ogun State, Nigeria.

⁴Department of Biochemistry, Ekiti State University, Ado-Ekiti, Nigeria.

⁵Department of Biochemistry, Federal University Oye Ekiti, Oye-Ekiti, Nigeria.

⁶Department of Biochemistry, Afe Babalola University, Ado-Ekiti, Nigeria.

*E-mail: ojo.adeleke@lmu.edu.ng

Abstract

Context: Cadmium has been considered as one of the most hazardous toxic compounds with harmful effect on the health of organisms.

Aims: To evaluate the effects of gallic acid (GA) on the cadmium-induced liver and renal oxidative stress in Wistar rats.

Methods: Twenty Wistar rats were grouped into four (A–D) of five rats. Rats in Group A, B, C and D were administered distilled water, 5 mg/kg bw cadmium chloride (CdCl₂), CdCl₂ + GA concurrently and GA (20 mg/kg bw) respectively and administered for 14 days. Biochemical parameters such as antioxidant enzyme activities, urea, creatinine and myeloperoxidase activity were determined.

Results: In the urea, creatinine and MPO, there was a significant increase in the CdCl₂ treated group. In the liver, the CdCl₂ treated group reduced significantly the catalase activity and increased the reduced glutathione. The gallic acid group increased in the GSH level, SOD, and CAT activities and it also reduced significantly the MDA level. However, the co-administration of CdCl₂ + GA had a considerably increase in the antioxidant enzymes. In the kidney, catalase activity and MDA level significantly decrease and increase respectively. The gallic acid also increases significantly the CAT and SOD activities while the MDA level was reduced. Co-administration of GA + CdCl₂ had a substantial increase only in the SOD activity compared to the control.

Conclusions: This study indicates that gallic acid was able to protect the alteration induced by cadmium chloride in the rat kidney and liver.

Keywords: gallic acid; cadmium chloride; antioxidant; nephrotoxicity; hepatotoxicity.

Resumen

Contexto: El cadmio ha sido considerado como uno de los compuestos tóxicos más peligrosos con efectos nocivos para la salud de los organismos.

Objetivos: Evaluar los efectos del ácido gálico (GA) sobre el estrés oxidativo renal y hepático inducido por cadmio en ratas Wistar.

Métodos: Se agruparon veinte ratas Wistar en cuatro (A–D) de cinco ratas. A las ratas de los grupos A, B, C y D se les administró agua destilada, 5 mg/kg de peso corporal de cloruro de cadmio (CdCl₂), CdCl₂ + GA al mismo tiempo y GA (20 mg/kg de peso corporal) respectivamente y durante 14 días. Se determinaron parámetros bioquímicos como actividad enzimática antioxidante, urea, creatinina y actividad mieloperoxidasa.

Resultados: En el grupo tratado con urea, creatinina y MPO, hubo un aumento significativo en el grupo tratado con CdCl₂. En el hígado, el grupo tratado con CdCl₂ redujo significativamente la actividad catalasa y aumentó el glutatión reducido. El grupo de ácido gálico aumentó en las actividades de nivel de GSH, SOD y CAT y también redujo significativamente el nivel de MDA. Sin embargo, la coadministración de CdCl₂ + GA tuvo un aumento considerable de las enzimas antioxidantes. En el riñón, la actividad de la catalasa y el nivel de MDA disminuyen y aumentan significativamente, respectivamente. El ácido gálico también aumenta significativamente las actividades de CAT y SOD mientras que se redujo el nivel de MDA. La coadministración de GA + CdCl₂ tuvo un aumento sustancial solo en la actividad de SOD en comparación con el control.

Conclusiones: Este estudio indica que el ácido gálico fue capaz de proteger la alteración inducida por el cloruro de cadmio en el riñón e hígado de rata.

Palabras Clave: ácido gálico; antioxidante; cloruro de cadmio; hepatotoxicidad; nefrotoxicidad.

ARTICLE INFO

Received: December 16, 2020.

Received in revised form: February 18, 2021.

Accepted: February 18, 2021.

Available Online: April 29, 2021.

AUTHOR INFO

ORCID: 0000-0001-9331-396X (OAO)



INTRODUCTION

The toxicity of heavy metals has become a public health challenge due to exposure to both occupational and environmental resources. This threat to environmental and public health is due to acute or chronic metal exposure brought about severe organ damage and even death (Ali et al., 2019). Cadmium (Cd) has been established as a known toxic substance abundantly present due to its importance in the industry worldwide. It is not biodegradable and has become a major health hazard as a result of its stretched half-life of about 10-30 years (Anetor et al., 2016). It is found in air, food, water, soil, and tobacco smoke. In occupational settings, its use involves Nickel-Cd battery production, zinc refining, smelting, welding, galvanizing, electroplating and by pigments, and plastics manufacturer (Odeyemi et al., 2011). At a lower concentration, Cd induces tissue damage. Cd has been considered as a category I cancer-causing agent by the International Agency for Research on Cancer (IARC, 1993). Cd toxicity has been characterized by proteinuria, bone fractures, severe pain, and osteomalacia, predominantly among women (Rahimzadeh et al., 2017).

Cd ingested is distributed via vital body part, for example, liver, lung, testis, kidney, heart, and brain, and most of it enters and accumulates into the liver and kidneys (Genchi et al., 2020). Cd causes cellular reactive oxygen species synthesis in the kidney by expanding lipid peroxidation. Acute and chronic exposure initiates adjacent tubular dysfunction and leads to nephrotoxicity. Nephrotoxicity induced structural damage of kidneys and clinically observed as aminoaciduria, glycosuria, and proteinuria (Ewere et al., 2016). The liver is one of the key vital organs for the harmfulness of Cd. It tends to be dependent upon particular clinical and morphological modifications under Cd impact. The hepatotoxicity of cadmium induces acute lethality in the body. There has been a strong relationship between the exposures to various environmental chemicals such as Cd and both renal and hepatic damage reported through sever-

al epidemiological studies (Rinaldi et al., 2017; Branca et al., 2018).

Several reports established that liver and kidney were the vast majority of delicate natural tissues readily influenced by Cd poisonousness (Ognjanovic et al., 2010; Ojo et al., 2014c). The initiation of Cd nephrotoxicity for instance generated through several pathways comprising radicals and cell death (El-Sharaky et al., 2007). Earlier studies focused on the use of metals for example zinc, copper, as antioxidants for defense against Cd toxicity. These inorganic ions were revealed to be involved in the incorporation and transference of Cd constituents inside biological environments via concealing and transferring processes (González-Trujano and Navarrete, 2011). Though, the usage of these ions in extreme portions might trigger severe biological complications.

Gallic acid (GA) is a naturally-occurring polyphenolic constituent present in fruits and several products of the soil plants. GA is an antioxidant found naturally in plants with application in the drug, food, and cosmetics industries (Owumi et al., 2020). GA possess several activities such as anti-mutagenic, antiviral, antifungal, antibacterial, against cancer-causing, anti-allergic, and anti-inflammatory ability through its antioxidant potentials (Olusoji et al., 2017; Zahrani et al., 2020). The scavenging ability of GA has been reported as a plausible component for the decrease of oxidative stress and improvement of degenerative disorders (Oyagbemi et al., 2016). GA also suppresses pro-inflammatory cytokines, reduces IL-6, and TNF- α expression suggesting it possesses anti-inflammatory properties (Mohamed et al., 2016). In this study, we evaluated the defensive and antioxidant impacts of GA against Cd-induced liver and renal stress in Wistar rats.

MATERIAL AND METHODS

Chemicals and reagents

Gallic acid (GA), cadmium chloride (CdCl₂), and also commercial assay kits of urea, creatinine,

and myeloperoxidase (MPO) were bought from Sigma-Aldrich Company (St. Louis, MO, USA).

Experimental animals and treatments

Twenty Wistar rats (170-200 g) were procured from the Department of Biochemistry, Landmark University. The animals were allowed to be acclimatized after which they were randomized into four groups (A-D) of five rats. The animals were exposed to light cycle (12h light/12h dark), room temperature of 24-27°C, and fed with commercial rat chow and water *ad libitum*. The experimental dealings on the rats complied with the ethical rules and affirmed by ethical committee (LUAC/2020/0052B) of the Department of Biochemistry, Landmark University, Nigeria.

Animal groupings

Rats in Group A were given distilled water daily, group B rats were orally given 5 mg/kg b.w. cadmium daily, group C rats were given cadmium (5 mg/kg b.w./daily) and GA (20 mg/kg b.w./day) concurrently, group D rats were administered GA (20 mg/kg b.w./daily). CdCl₂ and GA were dissolved in distilled water and administered for 2 weeks. Doses for cadmium and GA were selected based on earlier reports by El-Demerdash et al. (2004) for cadmium, and Ola-Davies and Olu-kole (2018) for gallic acid. It has been shown that the dose selected induces major oxidative stress in different tissues (Ojo et al., 2014a; 2014b; 2014c), while gallic acid was reported to be safe at 20 mg/kg body weight per day (Ola-Davies and Olu-kole, 2018).

Serum and organ preparation

The rats were anesthetized twenty-four hours after the last treatment under diethyl ether. The blood was collected, left for an hour, and the serum was separated by centrifuging at 1300×g for 10 min. The tissues of interest (liver and kidney) were removed, homogenized, and centrifuged at 5000 g for 10 minutes.

Biochemical parameters

The serum samples were utilized for estimation of kidney indices such as serum creatinine (CRE), urea, and myeloperoxidases (MPO) using commercial kits. The malondialdehyde (MDA) level, superoxide dismutase (SOD) and catalase (CAT), and the reduced glutathione (GSH) level were measured in the kidney and liver homogenates. The level of lipid peroxidation by measuring malondialdehyde level was according to Satoh (1978), CAT, SOD, and GSH content were determined according to Misra and Fridovich (1972), Aebi (1974), and Beutler (1963), respectively.

Histopathological study

For proper fixation, a portion of the liver and kidney tissues collected was immersed in 10% buffered formalin. These organs were inserted in paraffin wax using hematoxylin and eosin (H & E) staining, about 5–6 µm of the tissue were stained for histological assessment (Drury et al., 1976). Concisely, fixed liver and kidney tissues were dehydrated in methanol and xylene, respectively. Afterward, tissues sections (5 µm) were made, mounted on glass slides, and stained with stained with H&E. The slides were analyzed under the light microscope (Olympus BX63 with a DP72 camera, Olympus Corporation, Tokyo, Japan) at 400× and sections were observed.

Statistical analysis

Data were analyzed via one-way analysis of variance (ANOVA) along with Tukey's *posthoc* test on GraphPad Prism 8.0 (Version 8.0). Values were expressed as mean ± standard error of mean SEM (n = 5) with p significant at ≤0.05.

RESULTS

Table 1 revealed that CdCl₂ treated group produced considerable increment (p<0.05) in the concentrations of serum urea, creatinine, and MPO activities. Whereas, in the groups administered with GA alone and co-administration of CdCl₂ + GA showed a substantial reduction in the concentrations of urea, creatinine, and MPO activities.

Table 1. Protective impact of gallic acid on cadmium chloride-induced alterations on some selected serum markers.

Parameter	Control	Cadmium chloride treated rats	Cadmium chloride + gallic acid-treated rats	Gallic acid-treated rats
Urea ($\mu\text{mol/L}$)	22.34 \pm 1.11	32.89 \pm 1.42**	27.56 \pm 1.16*	21.56 \pm 1.07*
Creatinine ($\mu\text{mol/L}$)	0.44 \pm 0.02	0.62 \pm 0.08**	0.47 \pm 0.03*	0.49 \pm 0.04*
MPO ($\mu\text{mol/min}$)	22.45 \pm 1.56	45.62 \pm 6.43**	28.22 \pm 5.48*	21.76 \pm 1.89*


Values are expressed as mean \pm SEM (n=5). *p<0.05 significant difference compared to control (Group A); **p<0.05 statistically significant differences compared to CdCl₂ treated group. Myeloperoxidase (MPO); cadmium chloride (CdCl₂).

Fig. 1 displays the levels of enzymatic and non-enzymatic antioxidants in the liver. Hepatic SOD and CAT activities were significantly decreased in CdCl₂ treated groups compared to the control. Notwithstanding, SOD, CAT, and GSH levels improved substantially in the gallic acid-treated groups compared to the CdCl₂ group. Also, hepatic GSH concentrations was considerably elevated in the CdCl₂ group comparable to the control. In contrast, the considerable reduction was observed in the CdCl₂ + GA group. Hepatic MDA was expressively elevated in the CdCl₂ group compared to control, but considerably reduced in CdCl₂ + GA relative to the CdCl₂ treated group and gallic acid-treated groups relative to the control.

Fig. 2 reveals the concentrations of antioxidant enzymes and non-enzymatic antioxidants in the kidney. Renal SOD and CAT activities were altogether decreased in CdCl₂ groups compared to the control, but significantly increased in CdCl₂ + GA and GA treated groups relative to CdCl₂ treated and control groups. However, SOD and CAT activities were enhanced considerably in the GA groups compared to the control. Also, renal GSH

concentrations were slightly reduced in the CdCl₂ treated group compared to the control. Renal MDA was significantly increased in CdCl₂ group comparable to the control, but drastically reduced in the CdCl₂ + GA relative to CdCl₂ treated group and gallic acid groups relative to the control.

The defensive function of gallic acid against CdCl₂ toxicity was explored in hepatic and renal tissues via histo-architectural studies as revealed in Figs. 3-4. Normal physiology with no visible lesions was seen in the hepatocytes of control animals (Fig. 3) and gallic acid-treated group alone. In rats treated with CdCl₂ revealed mild vascular congestion while groups co-administered with CdCl₂ + gallic acid revealed regeneration. In the kidney histology, normal renal cortex, glomeruli, and tubules were seen in the control animals (Fig. 4) and gallic acid-treated group alone. In contrast, mild congestion of blood vessels and dilation of glomeruli were observed in CdCl₂-treated rats (Fig. 4). In groups co-administered with CdCl₂ + GA, revealed regeneration and restoration of the glomeruli and cells.








Figure 3. Cross-section of the liver of rats after two weeks. Group A (Control rats), Group B (cadmium chloride-treated rats), Group C (CdCl₂ + GA-treated rats), Group D (gallic acid-treated rats). (A) Control showing hepatocytes, normal portal tracts and central vein; (B) CdCl₂ group showing normal hepatocytes, normal portal tracts, and central vein and mild vascular congestion; (C) CdCl₂ + GA group showing mild vascular congestion; (D) GA group reveals normal hepatocytes, normal portal tracts, and central vein.



DISCUSSION

Cd exposure through contaminated food, water, air, commercial phosphate fertilizer, occupational hazards, and manufactured goods are a major sources of cadmium intoxication (Hayat et al., 2019). It was documented that contact with CdCl₂ mixtures exacerbates increment in the production of oxidants, which triggers oxidative injury of numerous natural organs (Liu et al., 2008). The liver and kidney have been described as the greatest delicate tissues to Cd toxicity (Liu et al., 2010; Ojo et al., 2014c). The build-up of CdCl₂ in human tissues contributes to several pathological disorders comprising hepatic and renal dysfunction (Ojo et al., 2014a; 2014c). So, evaluating the actions of poisonousness caused by heavy metals and medications urge more researchers to look for natural

products with essential antiradical property to shield biological organisms from oxidative injury caused by the generation of radicals (Sarwat et al., 2011). This investigation is meant to assess the effectiveness of gallic acid against cadmium-induced hepato-renal toxicity, via an antioxidant defense system.

In the current investigation, serum urea, creatinine, and MPO were evaluated to survey kidney performance. The information got indicated a huge increment in kidney indices; urea, MPO, and creatinine following CdCl₂ exposure. The alterations in the concentrations of serum kidney indices signify kidney injury trigger via CdCl₂ as earlier documented by (Purena et al., 2018). The increase in urea and creatinine might also be associated with hypertension-induced renal damage (Abdel-Zaher et al., 2019). These data correlate with other

researchers' work who documented that CdCl₂ initiated tube-shaped necrosis or loss of the membrane lining and injury of the kidney (Wang et al., 2009). In addition to that, the elevated serum urea produced by CdCl₂ toxicity might be identified with derangement in the catabolism of protein as an aftereffect of the rise in the production of the enzyme arginase implicated in urea generation (Tormanen, 2006). Furthermore, the administration of gallic acid revealed substantial enhancement in renal performance by a decline in the concentrations of urea and creatinine. Thus, the information acquired indicates a fundamental antioxidant property of gallic acid towards CdCl₂ toxicity, and that the treatment of gallic acid alone has no impact on the concentrations of kidney indices demonstrating the harmless usage of gallic acid on the liver and kidney.

Cd has been noted to stimulates the production of free radicals for example, hydroxyl radical (OH^\cdot), superoxide anion (O_2^\cdot), etc. In humans and animals, pathological conditions and oxidative deterioration of macromolecules are initiated by the overwhelming level of free radicals (Rani et al., 2014). Cd is absorbed and rapidly taken from the blood to the liver and kidney majorly, where it concentrates and induces many metabolic and histological changes, altered gene expression, apoptosis, and membrane damage (Matović et al., 2015; Andjelkovic et al., 2019). The efficient pathway of cellular defense and repair occurs via anti-oxidative protection against oxidant species. Antioxidant enzyme includes SOD and CAT. SOD catalyzes the reduction of O_2^\cdot to hydrogen peroxide (H_2O_2) whereas H_2O_2 is additionally separated via catalase to water and oxygen. Several reports in the literature have utilized different chelating agents to ease the harmfulness of Cd by increasing Cd excretion; nonetheless, chelation therapy is controversial in terms of efficiency and safety (Rahimzadeh et al., 2017; Rana et al., 2018; Ojo et al., 2018). Cd shows a high affinity for compounds containing the thiol group. In our study, lipid peroxidation increased with Cd administration while a decrease in catalase which establishes the toxicity already reported in the literature. The group administered with GA increased the antioxidant ac-

tivities and reduced the lipid peroxidation biomarker. However, the induced toxicity of cadmium was ameliorated by the administration of GA. This supports the claim of GA as an antioxidant agent. GSH is a nucleophilic substance containing sulfur with a high concentration in the liver and kidney. It promotes cellular protection from oxidative stress and other toxic compounds. Cd cytotoxicity has been linked to alteration in the metabolism of cellular GSH and its increase may protect cells from the induced Cd toxicity (Remelli et al., 2016). GA increases the intracellular GSH level by scavenging the reactive oxygen species (Hagar and Al Malki, 2014; Lee et al., 2018).

The build-up of Cd in rat liver caused organ damage (Ojo et al., 2014a). The histology of the hepatocytes in the control group shows normal portal tracts, and central vein while the GA group showing usual hepatocytes, portal tracts and central vein. The induced CdCl₂ group showed extensive vascular congestion. However, the CdCl₂ + GA group reveals mild vascular congestion. According to the histopathological changes seen in the kidneys, it may be concluded that cadmium harmfulness produced injury through developing tubular harm while the adjacent tubules were delicate to cadmium poisonousness due to their high re-absorptive activity (Ojo et al., 2014c).

CONCLUSIONS

This study indicates that gallic acid was able to protect the alteration induced by CdCl₂ in the rat kidney and liver by increasing the anti-oxidative enzymes and restore the histological changes induced by cadmium.

CONFLICT OF INTEREST

The authors declare no conflicts of interests.

ACKNOWLEDGMENTS

Authors acknowledge the technical support provided by the laboratory assistant. The authors acknowledge Landmark University Centre for Research, Innovation and Discovery (LUCRID) for the payment of the article charges. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

REFERENCES

- Abdel-Zaher AO, Abd-Ellatif RB, Aboulhagag NA, Farghaly HS, Al-Wasei FM (2019) The interrelationship between gasotransmitters and lead-induced renal toxicity in rats. *Toxicol Lett* 310: 39–50.
- Aebi H (1974) Catalase estimation. In: Bergmeyer HU, editor. *Methods of enzymatic analysis*. Weinheim (Germany)/New York (NY): Verlag Chemie/Academic Press Inc., p. 673–684.
- Ali H, Khan E, Ilahi I (2019) Environmental chemistry and ecotoxicology of hazardous heavy metals: environmental persistence, toxicity, and bioaccumulation. *J Chem* 2019: 6730305.
- Andjelkovic M, Buha Djordjevic A, Antonijevic E, Antonijevic B, Stanic M, Kotur-Stevuljevic J, Spasojevic-Kalimanovska V, Jovanovic M, Boricic N, Wallace D, Bulat Z (2019) Toxic effect of acute cadmium and lead exposure in rat blood, liver, and kidney. *Int J Environ Res Public Health* 16(2): 274.
- Anetor JI, Uche CZ, Ayita EB, Adedapo SK, Adeleye JO, Anetor GO, Akinlade SK (2016) Cadmium level, glycemic control, and indices of renal function in treated type II diabetics: implications for polluted environments. *Front Public Health* 13(4): 114.
- Beutler E, Durgun O, Kelly BM (1963) Improved method for the determination of blood glutathione. *J Lab Clin Med* 51: 882–888.
- Branca JJ, Morucci G, Pacini A (2018) Cadmium-induced neurotoxicity: still much ado. *Neural Regen Res* 13(11): 1879.
- Drury RA, Wallington EA, Cancerson R (1976) *Carlton's Histopathological Techniques* (4th ed.). Oxford, UK: Oxford University Press
- El-Demerdash FM, Yousef MI, Kedwany FS, Baghdadi HH (2004) Cadmium-induced changes in lipid peroxidation, blood hematology, biochemical parameters and semen quality of male rats: protective role of vitamin E and β -carotene. *Food Chem Toxicol* 42(10): 1563e1571.
- El-Sharaky AS, Newairy AA, Badreldeen MM, Eweda SM, Sheweita SA (2007) Protective role of selenium against renal toxicity induced by cadmium in rats. *Toxicology* 235: 185–193.
- Ewere EG, Oyebadejo SA, Peter VC (2016) Ethanolic leaf extract of *Iringia gabonensis* (O'Rorke) Baill protects against nephrotoxicity and hepatotoxicity in cadmium-induced Wistar albino rats. *Int J Pharm Toxicol* 4(2): 105.
- Genchi G, Sinicropi MS, Lauria G, Carocci A, Catalano A (2020) The effects of cadmium toxicity. *Int J Environ Res Public Health* 17(11): 3782.
- González-Trujano E, Navarrete A (2011) Effect of zinc on the cadmium acute intoxication in the gastric injury induced in rats. *Rev Latinoamer Quím* 39 (1–2): 45–54.
- Hagar H, Al Malki W (2014) Betaine supplementation protects against renal injury induced by cadmium intoxication in rats: role of oxidative stress and caspase-3. *Environ Toxicol Pharmacol* 37(2): 803–811.
- Hayat MT, Nauman M, Nazir N, Ali S, Bangash N (2019) Environmental hazards of cadmium: past, present, and future. In *Cadmium Toxicity and Tolerance in Plants*; (pp. 163–183). Academic Press.
- IARC (1993) Working Group on the Evaluation of Carcinogenic Risks to Humans. Beryllium, cadmium, mercury, and exposures in the glass manufacturing industry. International Agency for Research on Cancer.
- Lee S, Li J, Zhou X, Yin J, Yoon J (2018) Recent progress on the development of glutathione (GSH) selective fluorescent and colorimetric probes. *Coord Chem Rev* 366: 29–68.
- Liu J, Qian SY, Guo Q, Jiang J, Waalkes MP, Mason RP, Kadiiska MB (2008) Cadmium generates reactive oxygen- and carbon-centered radical species in rats: insights from *in vivo* spin-trapping studies. *Free Radic Biol Med* 45: 475–481.
- Liu J, Huang HL, Zhang WC, Li H (2010) Cadmium-induced increase in uterine wet weight and its mechanism. *Birth Defects Res B Dev Reprod Toxicol* 89: 43–49.
- Matović V, Buha A, Đukić-Čosić D, Bulat Z (2015) Insight into the oxidative stress induced by lead and/or cadmium in blood, liver and kidneys. *Food Chem Toxicol* 78: 130–140.
- Misra HP, Fridovich I (1972) The role of superoxide anion in the autooxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem* 247: 3170–3175.
- Mohamed HM, Abd El-Twab SM (2016) Gallic acid attenuates chromium-induced thyroid dysfunction by modulating antioxidant status and inflammatory cytokines. *Environ Toxicol Pharmacol* 48: 225–236.
- Odewumi CO, Badisa VL, Le UT, Latinwo LM, Ikediobi CO, Badisa RB, Darling-Reed SF (2011) Protective effects of N-acetylcysteine against cadmium-induced damage in cultured rat normal liver cells. *Int J Mol Med* 27(2): 243–248.
- Ognjanović BI, Marković SD, Ethordević NZ, Trbojević IS, Stajn AS, Sačić ZS (2010) Cadmium-induced lipid peroxidation and changes in antioxidant defense system in the rat testes: protective role of coenzyme Q (10) and vitamin E. *Reprod Toxicol* 29: 191–197.
- Ojo OA, Ajiboye BO, Oyinloye BE, Ojo AB (2014a) Prophylactic effects of ethanolic extract of *Iringia gabonensis* stem bark against cadmium-induced toxicity in albino rats. *Adv Pharm* 8: 894610.
- Ojo OA, Oyinloye BE, Ajiboye BO, Onikanni SA (2014b) Neuroprotective mechanism of ethanolic extract of *Iringia gabonensis* stem bark against cadmium-induced neurotoxicity in rats. *Br J Med Med Res* 4(36): 5793–5805.
- Ojo OA, Ajiboye BO, Oyinloye BE, Ojo AB, Olarewaju OI (2014c) Protective effect of *Iringia gabonensis* stem bark extract on cadmium-induced nephrotoxicity in rats. *Interdiscip Toxicol* 7(4): 208–214.

- Ojo OA, Ojo AB, Awoyinka OA, Olayide I, Ajiboye BO, Oyinloye BE, Osukoya OA, Ibitayo AO (2018) Aqueous extract of *Carica papaya* Linn roots potentially attenuates arsenic induced biochemical and genotoxic effects in Wistar rats. *J Tradit Complement Med* 8(2): 324–334.
- Ola-Davies OE, Olukole SG (2018) Gallic acid protects against bisphenol A-induced alterations in the cardiorenal system of Wistar rats through the antioxidant defense mechanism. *Biomed Pharmacother* 107: 1786–1794.
- Olusoji MJ, Oyeyemi OM, Asenuga ER, Omobowale TO, Ajayi OL, Oyagbemi AA (2017) Protective effect of gallic acid on doxorubicin-induced testicular and epididymal toxicity. *Andrologia* 49(4): e12635.
- Owumi S, Najophe ES, Farombi EO, Oyelere AK (2020) Gallic acid protects against Aflatoxin B1-induced oxidative and inflammatory stress damage in rats kidneys and liver. *J Food Biochem* 4: e13316.
- Oyagbemi AA, Omobowale TO, Saba AB, Olowu ER, Dada RO, Akinrinde AS (2016) Gallic acid ameliorates cyclophosphamide-induced neurotoxicity in Wistar rats through free radical scavenging activity and improvement in antioxidant defense system. *J Diet Suppl* 13(4): 402–419.
- Purena R, Seth R, Bhatt R (2018) Protective role of *Emblica officinalis* hydro-ethanolic leaf extract in cisplatin induced nephrotoxicity in rats. *Toxicol Rep* 5: 270–277.
- Rahimzadeh MR, Rahimzadeh MR, Kazemi S, Moghadamnia AA (2017) Cadmium toxicity and treatment: An update. *Caspian J Intern Med* 8(3): 135.
- Rana MN, Tangpong J, Rahman MM (2018) Toxicodynamics of lead, cadmium, mercury and arsenic-induced kidney toxicity and treatment strategy: a mini review. *Toxicol Rep* 5: 704–713.
- Rani A, Kumar A, Lal A, Pant M (2014) Cellular mechanisms of cadmium-induced toxicity: a review. *Int J Environ Health Res* 24(4): 378–399.
- Remelli M, Nurchi VM, Lachowicz JI, Medici S, Zoroddu MA, Peana M (2016) Competition between Cd (II) and other divalent transition metal ions during complex formation with amino acids, peptides, and chelating agents. *Coord Chem Rev* 327: 55–69.
- Rinaldi M, Micali A, Marini H, Adamo EB, Puzzolo D, Pisani A, Trichilo V, Altavilla D, Squadrito F, Minutoli L (2017) Cadmium, organ toxicity and therapeutic approaches: a review on brain, kidney and testis damage. *Curr Med Chem* 24(35): 3879–3893.
- Sarwat M, Das S, Srivastava PS (2011) Estimation of genetic diversity and evaluation of relatedness through molecular markers among medicinally important trees: *Terminalia arjuna*, *T. chebula* and *T. bellerica*. *Mol Biol Rep* 38: 5025–5036.
- Satoh K (1978) Serum lipid peroxide in cerebrovascular disorders determined by a new colorimetric method. *Clin Chim Acta* 90: 37–43.
- Tormanen CD (2006) Inhibition of rat liver and kidney arginase by cadmium ion. *J Enzyme Inhib Med Chem* 21: 119–123.
- Wang L, Chen D, Cao J, Liu Z (2009) Protective effect of N-acetylcysteine on experimental chronic cadmium nephrotoxicity in immature female rats. *Hum Exp Toxicol* 28: 221–229.
- Zahrani NA, El-Shishtawy RM, Asiri AM (2020) Recent developments of gallic acid derivatives and their hybrids in medicinal chemistry: A review. *Eur J Med Chem* 20: 112609.

AUTHOR CONTRIBUTION:

Contribution	Ojo OA	Rotimi D	Bright AE	Kayode O	Ojo AB	Alejolowo O	Ajiboye BO	Oluba OM
Concepts or ideas	x				x			x
Design	x	x		x		x	x	x
Definition of intellectual content	x	x		x	x		x	x
Literature search		x	x		x	x		
Experimental studies		x	x					
Data acquisition		x	x	x	x	x		
Data analysis	x						x	x
Statistical analysis		x	x			x		
Manuscript preparation	x	x		x	x	x		
Manuscript editing	x			x	x	x	x	x
Manuscript review	x	x	x	x	x	x	x	x

Citation Format: Ojo OA, Rotimi D, Bright AE, Kayode O, Ojo AB, Alejolowo O, Ajiboye BO, Oluba OM (2021) Gallic acid protects against cadmium chloride-induced alterations in Wistar rats via the antioxidant defense mechanism. *J Pharm Pharmacogn Res* 9(5): 668–676.